

Name _____ DOB _____

TUBERCULOSIS SCREENING FORM

A Tuberculosis skin test is required within three months prior to admission at St. John's College of **ALL F1/INTERNATIONAL STUDENTS AND ANY OTHER US CITIZEN WHO HAS TRAVELED OUTSIDE OF THE USA WITHING THE PAST 5 YEARS FOR 30 DAYS OR MORE.** If the student has had a BCG vaccine, a blood test (Quantiferon TB Gold) is required in place of the skin test. The college requirement for tuberculosis screening is consistent with the recommendations of the U.S. Centers for Disease Control and Protection and with the American College Health Association's Guidelines for Tuberculosis Screening.

1. Tuberculin Skin Test (PPD)

Date Given (Mo./Day/Yr.)	Date Read (Mo./Day/Yr.)	Result (record actual mm of induration; transverse diameter; if no induration, write "0")	Interpretation** (based on mm of induration and risk factors)
		_____mm	Positive/Negative

2. If Tuberculin Skin Test is positive, chest x-ray required**

Date of chest x-ray (Mo./Day/Yr.)	Result (please circle)	Attach copy of chest x-ray report (required)
	Normal/Abnormal	

3. Document any treatment patient has received for either a positive TB skin test or active tuberculosis.

Medication(s) received _____

Date started (Mo./Day/Yr.)	Date Completed (Mo./Day/Yr.)

4. Health Care Provider Name, Contact Information, and Signature

****Interpretation guidelines**

>5 mm is positive:

- Recent close contacts of an individual with infectious TB
- Persons with fibrotic changes on a prior chest x-ray consistent with past TB disease
- Organ transplant recipients
- Immunosuppressed persons: taking > 15 mg/d of prednisone for > 1 month; taking a TNF- α antagonist
- Persons with HIV/AIDS

>10 mm is positive:

- Persons born in a high prevalence country or who resided in one for a significant* amount of time
- History of illicit drug use
- Mycobacteriology laboratory personnel
- History of resident, worker or volunteer in high-risk congregate settings
- Persons with the following clinical conditions: silicosis, diabetes mellitus, chronic renal failure, leukemias and lymphomas, head, neck or lung cancer, low body weight (>10% below ideal), gastrectomy or intestinal bypass, chronic malabsorption syndromes

> 15 mm is positive

- Persons with no known risk factors for TB disease